

Cerebrofacial Venous Anomalies, Sinus Pericranii, Ocular Abnormalities and Developmental Delay

B. MACIT¹, P.E. BURROWS², S. YILMAZ^{1,3}, D.B. ORBACH^{1,3}, J.B. MULLIKEN^{3,4}, A.I. ALOMARI^{1,3}

¹ Division of Vascular and Interventional Radiology, Children's Hospital Boston and Harvard Medical School; Boston, MA, USA

² Department of Diagnostic Imaging, Texas Children's Hospital; Houston, TX, USA

³ Vascular Anomalies Center; ⁴ Department of Plastic Surgery, Children's Hospital Boston and Harvard Medical School; Boston, MA, USA

Key words: sinus pericranii, seizure disorder, venous malformation, developmental delay

Summary

The clinical implications of venous cerebrovascular maldevelopment remain poorly understood. We report on the association of cerebrofacial venous anomalies (including sinus pericranii), ocular abnormalities and mild developmental delay in two children. In addition, one child had a seizure disorder. Complex cerebrofacial slow-flow vascular anomalies may herald an underlying developmental aberration affecting the cerebrofacial and orbital regions.

Introduction

The coexistence of intracranial and extracranial vascular anomalies has been well documented in the literature ¹⁻³. Abnormal development of the brain has also been associated with anomalies of the cerebral and facial vessels ⁴⁻⁷. Sinus pericranii refers to an abnormal communication between the intracranial dural venous sinuses and anomalous epicranial veins. In this communication, we describe two patients with cerebrofacial venous anomalies including sinus pericranii, in association with ocular abnormalities, mild developmental delay and possibly seizures.

Case Reports

Institutional review board approval was not required by our hospital for publishing retrospective case reports. However, in keeping with the ethical conduct of studies, the principles of the Declaration of Helsinki were followed.

First Patient

An eight-year-old boy was referred to our center for treatment of cerebrofacial vascular anomalies. The patient was born with a vascular "birthmark" on the forehead and around the left eye, and he had a medically-controlled seizure disorder. Examination showed a small left orbit and globe, with compromised visual acuity, proptosis, ptosis, glaucoma, miosis and dense retropupillary membranes (Figure 1A). There was a palpable bony defect in the forehead with overlying compressible, non-pulsatile bluish soft tissue which distended in the dependent position. The child had academic difficulties in reading and writing, with attention deficit disorder.

Cross-sectional studies demonstrated a smaller left cerebral hemisphere, with focal calcification and a cavernous malformation in the basal ganglia (Figure 1B,C). The venous malformation in the left orbit and frontal region




Figure 1 A) First Patient. There is a small left orbit and globe with venous expansion of the left forehead. B) Non-contrast enhanced head CT scan reveals a focal calcification in the left basal ganglia (small arrow) and venous malformation of the scalp (large arrow). C) Enhanced Axial T1-weighted MR Image shows smaller left cerebral hemisphere with DVAs in the right caudate nucleus and thalamus (small arrow). The anomalous transcalvarial and epicranial veins are clearly depicted (large arrow). D) Venous phase of cerebral angiography shows global anomalies of the venous drainage of the brain. Note the presence of persistent falcine sinus (long arrow) and fenestrated, underdeveloped superior sagittal sinus with retrograde filling of the sinus pericranii (short arrows).




Figure 2 A) Second Patient. An overgrown left cheek with venous malformation extending into the upper lip and upper eyelid. B) Sonographic image of the left orbit demonstrates linear echogenic septa (arrow) within the posterior aspect of the vitreous space. C) Venous phase of cerebral angiography show transcalvarial communication between the superior sagittal sinus and the epicranial venous malformation.

traversing the left frontal calvaria represents a sinus pericranii. There were multiple complex developmental venous anomalies (DVA) in the left cerebral hemisphere. Cerebral angiography demonstrated several anomalies of the venous drainage of both cerebral hemispheres, including the presence of persistent falcine sinus and fenestrated, underdeveloped superior sagittal sinus with retrograde filling of the sinus pericranii.

Second Patient

A three-year-old boy was born with a vascular "birthmark" on the left cheek that progressively enlarged. A small bony groove was felt on the left side of the calvaria at the age of one year. The child had minor developmental delay

with attention deficit disorder, learning disabilities, and delayed speech. There was a reticular blue stain overlying an enlarged left cheek, upper eyelid and oral mucosa. The cheek was soft and compressible. He had an irregular groove on the left parietal calvaria with large overlying venous channels.

MRI study showed a venous malformation of the left parietal calvaria with sinus pericranii extending into the left upper eyelid and masseter muscle (Figure 2A) with a left cerebral DVA. An ultrasonographic examination of the left orbit revealed linear echogenic membranes within the vitreous space with no significant flow within these membranes (Figure 2B). The visual acuity of the left eye was decreased, with ptosis and no retinal detachment. Cerebral an-

giography demonstrated normal arterial and cerebral venous anatomy with opacification of the sinus pericranii connecting the superior sagittal sinus through a parietal calvarial defect to anomalous tortuous scalp veins (Figure 2C).

Discussion

Sinus pericranii connects the extracranial and intracranial venous channels, typically in the midline frontal region⁷⁻⁸. Sinus pericranii is usually asymptomatic, although local pain, headache, nausea, and vertigo have been reported⁹.

Sinus pericranii has been reported in association with variable slow-flow vascular anomalies of the cerebrofacial region, including cerebral and cerebellar cavernous malformation, cerebral DVAs, venous malformations of the tongue and blue rubber bleb nevus syndrome^{7,10-12}. Konez et al. reported sinus pericranii in six out of 53 patients with cervicofacial venous malformation¹³.

In addition to the cerebrofacial venous anomalies, both patients presented have mild developmental delay and one seizure disorder. Different forms of cerebrofacial vascular anomalies have been described in association with developmental delay including sinus pericranii, dural sinus malformation, cerebral “angiomas” and anomalous cerebral veins¹⁴⁻¹⁶.

The coexistence of intracranial and extracranial vascular anomalies is not rare. Developmental venous anomalies are present in 20% of patients with extensive cervicofacial venous malformation¹⁷. Forty-five percent of patients with periorbital lymphatic malformation were found to have a cerebral DVA²⁻³. Bisdorf et al.³ reported that 70% of orbital lymphatic malformation had intracranial vascular anomalies. Cerebrofacial venous metameric syndromes

can be categorized into three main types: CVMS 1 (the medial prosencephalic or olfactory group with forehead and nose involvement), CVMS 2 (the lateral prosencephalic or optic group with involvement of the occipital lobe, eye, cheek, and maxilla) and CVMS 3 (the rhombencephalon or otic group with involvement of the cerebellum, lower face and mandible)¹⁸⁻¹⁹. Wyburn-Mason (or Bonnet-Dechaume-Blanc) syndrome is characterized by arteriovenous malformation of the brain, the orbit and the maxillofacial region¹. The major features of Sturge-Weber syndrome are facial capillary stain, leptomeningeal and choroidal small vessel anomalies, mental retardation, seizure, weakness, visual impairment and glaucoma²⁰. Bannayan-Riley-Ruvalcaba syndrome combines vascular anomalies (such as cerebral venous anomalies and fast-flow lesions) with variable degrees of developmental delay²¹.

Vascular insults are implicated in abnormal cerebral development⁴⁻⁶. Barkovich⁴ reported four patients with focal migrational anomalies associated with large vascular channels, three of them had developmental delay with seizure disorder. There are several case reports of the association of cortical migration disorder with arterial anomalies²²⁻²⁵. The association of developmental venous anomaly and neuromigrational disorders was reported by Riel-Romero et al.²⁵. An early embryonic vascular aberration affecting the cervicofacial endothelium (mesodermal) or the tunica media (from the neural crest) may lead to migration anomalies with segmental distribution in the regions of the face and brain¹.

In conclusion, the combination of cerebrofacial vascular anomalies with sinus pericranii, ocular abnormalities, developmental delay and possibly seizure disorder might represent a rare association that has not been previously appreciated.

References

- 1 Krings T, Geibprasert S, Luo CB, et al. Segmental neurovascular syndromes in children. *Neuroimag Clin N Am*. 2007; 17: 245-258.
- 2 Greene AK, Burrows PE, Smith L, et al. Periorbital lymphatic malformation: Clinical course and management in 42 patients. *Plast Reconstr Surg*. 2005; 115: 22-30.
- 3 Bisdorff A, Mulliken JB, Carrico J, et al. Intracranial vascular anomalies in patients with periorbital lymphatic and lymphaticovenous malformations. *Am J Neuroradiol*. 2007; 28: 335-341.
- 4 Barkovich JA. Abnormal vascular drainage in anomalies of neuronal migration. *Am J Neuroradiol*. 1988; 9: 939-942.
- 5 Stewart RM, Richman DP, Caviness VS Jr. Lissencephaly and pachygyria. An architectonic and topographical analysis. *J Neuropathol Exp Neurol*. 1974; 31: 1-12.
- 6 Richman DP, Stewart RM, Caviness VS Jr. Cerebral microgyria in a 27-week fetus: an architectonic and topographical analysis. *Acta Neuropathol*. 1974; 33: 374-384.
- 7 Gandolfo C, Krings T, Alvarez H, et al. Sinus pericranii: Diagnostic and therapeutic considerations in 15 patients. *Neuroradiology*. 2007; 49: 505-514.
- 8 Luker GD, Siegel MJ. Sinus pericranii: Sonographic findings. *Am J Roentgenol*. 1995; 165: 175-176.
- 9 Anegawa S, Hayashi T, Torigoe R, et al. Sinus pericranii with severe symptom due to transient disorder of venous return - Case report. *Neurol Med Chir (Tokyo)*. 1991; 31: 287-289.
- 10 Desai K, Bhayani R, Goel A, et al. Sinus pericranii in the frontal region: A case report. *Neurol India*. 2001; 49: 305-307.
- 11 Sakai K, Namba K, Meguro T, et al. Sinus pericranii associated with a cerebellar venous angioma - Case report. *Neurol Med Chir (Tokyo)*. 1997; 37: 464-467.
- 12 Sherry RG, Walker ML, Olds MV. Sinus pericranii and venous angioma in the blue-rubber bleb nevus syndrome. *Am J Neuroradiol*. 1984; 5: 832-834.
- 13 Konez O, Burrows PE, Mulliken JB. Cervicofacial venous malformations. MRI features and interventional strategies. *Interv Neuroradiol*. 2002; 8: 227-234.
- 14 Barbosa M, Mahadevan J, Weon YC, et al. Dural sinus malformations with giant lakes, in neonates and infants. *Interv Neuroradiol*. 2003; 9: 407-424.
- 15 Ficcadenti A, Santoro L, Petroni V, et al. Mental retardation, facial anomalies, brachydactyly, cerebral angiomas, femoral nucleus necrosis: A new entity or Hall-Riggs Syndrome? *Am J Med Genet A*. 2009; 149A: 1001-1005.
- 16 Nakayama T, Matsukado Y. Sinus pericranii with aneurismal malformation of internal cerebral vein. *Surg Neurol*. 1975; 3: 133-137.
- 17 Boukobza M, Enjolras O, Guichard JP, et al. Cerebral developmental venous anomalies associated with head and neck venous malformations. *Am J Neuroradiol*. 1996; 17: 987-994.
- 18 Ramli N, Sachet M, Bao C, et al. Cerebrofacial venous metamerism syndrome (CVMS) 3: Sturge-Weber syndrome with bilateral lymphatic/venous malformations of the mandible. *Neuroradiology*. 2003; 45: 687-690.
- 19 Agid R, Terbrugge KG. Cerebrofacial venous metamerism syndrome 2 plus 3: facial and cerebral manifestations. *Interv Neuroradiol*. 2007; 13: 55-58.
- 20 Thomas-Sohl KA, Vaslow DF, Maria BL. Sturge-Weber syndrome: a review. *Pediatr Neurol*. 2004; 30: 303-310.
- 21 Tan WH, Baris HN, Burrows PE, et al. The spectrum of vascular anomalies in patients with PTEN mutations: implications for diagnosis and management. *J Med Genet*. 2007; 44: 594-602.
- 22 Shankar JJ, Banerjee ST, Hogan M, et al. A rare case of cerebral cortical dysplasia with arterial vascular dysplasia. *Can J Neurol Sci*. 2009; 36: 757-760.
- 23 Araki Y, Mori S, Kanoh M, et al. Congenital hemicerebral arterial ectasia complicating unilateral megalencephaly. *Br J Radiol*. 1987; 60: 395-400.
- 24 Doran SE, Deveikis JP, and Chandler WF. Dolichoectasia of the anterior cerebral arteries in an adolescent. *Am J Neuroradiol*. 1975; 16: 1548-1550.
- 25 Abe T, Singer RJ, Marks MP, et al. Arterial vascular abnormality accompanying cerebral cortical dysplasia. *Am J Neuroradiol*. 1997; 18: 144-146.
- 26 Riel-Romero RMS, Mattingly M. Developmental venous anomaly in association with neuromigrational anomalies. *Pediatr Neurol*. 2005; 32: 53-55.

Ahmad I. Alomari, MD
 Division of Vascular and Interventional Radiology
 Children's Hospital Boston,
 300 Longwood Avenue
 Boston, MA 02115
 Tel.: +1-617-355-6221
 Fax: +1-617-730-0541
 E-mail: ahmad.alomari@childrens.harvard.edu